My dosing card for Hemlibra (emicizumab)

STARTING DOSE

3 mg/kg once a week for the first 4 weeks

Date:/	//	My weight:	kg
My dose:			
	AMOUN	T TO USE:	
Sky Blue*	Purple	Turquoise	Brown
30 mg/1 mL	60 mg/0.4 mL	105 mg/0.7 mL	150 mg/1 mL
(30 mg/mL)	(150 ma/mL)	(150 mg/mL)	(150 ma/mL)

ONGOING DOSE From week 5 Date: ____ /___ My weight: _ 1.5 mg/kg every week My dose: (tick correct box) 3.0 mg/kg every 2 weeks 6.0 ma/ka every 4 weeks Total dose: **AMOUNT TO USE:** Sky Blue* **Turquoise Purple Brown** 30 mg/1 mL 60 mg/0.4 mL 105 mg/0.7 mL 150 mg/1 mL (150 mg/mL) (150 mg/mL) (150 mg/mL)





Each colour relates to the different caps used on each vial strength of Hemlibra

*Do not combine Sky Blue topped vials containing 30 mg/1 ml of emicizumab with any other colour vial (Purple, Turquoise or Brown) containing 150 mg/ml emicizumab in the same injection due to the different concentrations of emicizumab between vials

(30 mg/mL)

My injection day will be:	

Fill in the date and tick the orange box when you complete each of your weekly injections.

STARTING DOSE		
DATE	INJECTION(S) COMPLETED	

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. You should also report side effects to Roche Products Ltd by emailing the Roche Drug Safety Centre at welwyn.uk_dsc@roche.com or calling +44 (0) 1707 367554. By reporting side effects you can help provide more information on the safety of this medicine.

ONGOING DOSE INJECTION(S) DATE **COMPLETED**